



Available at
WWW.MATHEMATICSWEB.ORG
 POWERED BY SCIENCE @ DIRECT®

Journal of
**MATHEMATICAL
 ANALYSIS AND
 APPLICATIONS**

J. Math. Anal. Appl. 282 (2003) 56–62

www.elsevier.com/locate/jmaa

A generalization of discrete Gronwall inequality and its application to weakly singular Volterra integral equation of the second kind

Lü Tao ^{a,*} and Huang Yong ^{b,c}

^a *Institute of Mathematics, Sichuan University, Chengdu 610064, PR China*

^b *Institute of Computer Application, Academia Sinica, Chengdu 610041, PR China*

^c *Institute of Business and Administration, Sichuan University, Chengdu 610064, PR China*

Received 13 December 2001

Submitted by R.P. Agarwal

Abstract

This paper presents a new discrete Gronwall inequality. Using the inequality, we prove convergence and error estimate of the numerical solutions of the second weakly singular Volterra integral equation, where discrete equation is derived by Novot's quadrature formula.
 © 2003 Elsevier Science (USA). All rights reserved.

1. Introduction

If a nonnegative sequence $\{y_n, n = 0, \dots, N\}$ satisfies

$$y_0 = 0, \quad y_n \leq A + Bh \sum_{j=0}^{n-1} y_j, \quad 1 \leq n \leq N, \quad h = 1/N, \quad (1)$$

then

$$\max_{0 \leq i \leq N} y_i \leq Ae^B, \quad (2)$$

where A and B are positive constants independent of h . As a discrete analogue of Gronwall inequality [5] this inequality is called discrete Gronwall inequality, which plays

* Corresponding author.

E-mail address: lutao@public.cd.sc.cn (L. Tao).

an important role in proving convergence of numerical solutions of Volterra integral equations with a continuous kernel [3,4,9]. But it is ineffective to prove convergence of numerical solutions of Volterra integral equations of the second kind with a weakly singular kernel. For example, we consider the following nonlinear Volterra equation of the second kind

$$u(s) = y(s) + \int_a^s k^*(s, t, u(t)) dt, \quad a \leq s \leq b, \quad (3)$$

where the kernel

$$k^*(s, t, u(t)) = (s - t)^\alpha (\ln |s - t|)^\beta k(s, t, u(t)), \quad -1 < \alpha \leq 0, \beta = 0, 1, \quad (4)$$

is weakly singular and $k(s, t, u(t))$ is a continuous function on variables s, t, u , especially, there exists a positive constant L satisfying

$$|k(s, t, u) - k(s, t, v)| \leq L|u - v|, \quad a \leq t, s \leq b. \quad (5)$$

Moreover for fixed s and t , we assume that $k(s, t, u(t))$ has high order derivatives on u , and let $k_u(s, t, u(t)) = \frac{\partial}{\partial u} k(s, t, u(t))$. In order to get a discrete equation of (3), we can apply Navot's quadrature formula of computing integrals with the end point singularity. Consider the integral

$$I(G) = \int_a^b G(x) dx = \int_a^b (b - x)^\alpha (\ln |b - x|)^\beta g(x) dx. \quad (6)$$

where $-1 < \alpha < 0$, $\beta = 0, 1$, and $G(x) = (b - x)^\alpha (\ln |b - x|)^\beta g(x)$, $g(x)$ is smooth on $[a, b]$. Take the step width $h = (b - a)/N$, and $x_i = a + ih$, $i = 0, \dots, N$. If $g(x) \in C^{2m}[a, b]$, then Navot [7] and Lyness [6] proved that the quadrature formula

$$Q_N(G) = \frac{h}{2} G(x_0) + h \sum_{i=1}^{N-1} G(x_i) - [-\beta \zeta'(-\alpha) + \zeta(-\alpha)(\ln h)^\beta] g(b) h^{1+\alpha} \quad (7)$$

possesses the following Euler–Maclaurin asymptotic expansion

$$\begin{aligned} E_N(G) = Q_N(G) - I(G) &= \sum_{j=1}^{m-1} \frac{B_{2j}}{(2j)!} G^{(2j-1)}(a) h^{2j} \\ &\quad + \sum_{j=1}^{2m-1} (-1)^j [-\beta \zeta'(-\alpha - j) + \zeta(-\alpha - j)(\ln h)^\beta] \\ &\quad \times g^{(j)}(b) h^{j+\alpha+1} + O(h^{2m}), \end{aligned} \quad (8)$$

where B_{2j} are Bernoulli numbers, and $\zeta(x)$ is the Riemann Zeta function. From (8) we know that

$$E_N(G) = O(h^{2+\alpha} |\ln h|^\beta), \quad (9)$$

if $g(x) \in C^2[a, b]$. Taking $s = x_i$ in (3) and using the quadrature formula (7) for

$$u(x_i) = y(x_i) + \int_{x_0}^{x_i} (x_i - t)^\alpha (\ln |x_i - t|)^\beta k(x, t, u(t)) dt, \quad (10)$$

we obtain the following nonlinear discrete equations: find $u_i, i = 0, 1, \dots, N$, satisfying

$$\begin{cases} u_0 = y(x_0), \\ u_i = y(x_i) + \frac{h}{2}(x_i - x_0)^\alpha (\ln |x_i - x_0|)^\beta k(x_i, x_0, u_0) \\ \quad + h \sum_{j=1}^{i-1} (x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta k(x_i, x_j, u_j) \\ \quad - [-\beta \zeta'(-\alpha) + \zeta(-\alpha)(\ln h)^\beta] k(x_i, x_i, u_i) h^{1+\alpha}, \quad i = 1, \dots, N. \end{cases} \quad (11)$$

But by means of (8) the integral equation (3) can be expressed as follows

$$\begin{aligned} u(x_i) = & y(x_i) + h \sum_{j=0}^{i-1} w_{ij} (x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta k(x_i, x_j, u(x_j)) \\ & + h w_{ii} k(x_i, x_i, u(x_i)) + E_{i,t}((x_i - t)^\alpha (\ln |x_i - t|)^\beta k(x_i, t, u(t))), \\ & i = 0, 1, \dots, N, \end{aligned} \quad (12)$$

where

$$\begin{aligned} w_{i0} &= \frac{1}{2}, \quad w_{ii} = h^\alpha [\beta \zeta'(-\alpha) - \zeta(-\alpha)(\ln h)^\beta], \\ \text{and } w_{ij} &= 1, \quad \text{for } 1 \leq j < i, \end{aligned} \quad (13)$$

and the remainder has an estimation

$$|E_{i,t}((x_i - t)^\alpha (\ln |x_i - t|)^\beta k(x_i, t, u(t)))| = O(h^{2+\alpha} |\ln h|^\beta). \quad (14)$$

Letting $e_i = u(x_i) - u_i$ and subtracting (11) from (12), we obtain the error $\{e_i\}$ satisfies the equation

$$\begin{cases} e_0 = 0, \\ e_i = h \sum_{j=0}^{i-1} w_{ij} (x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta [k(x_i, x_j, u(x_j)) - k(x_i, x_j, u_j)] \\ \quad + h w_{ii} [k(x_i, x_i, u(x_i)) - k(x_i, x_i, u_i)] \\ \quad + E_{i,t}(x_i, t, u(t)), \quad 1 \leq i \leq N. \end{cases} \quad (15)$$

From (5) we get

$$\begin{aligned} |e_i| &\leq Lh \sum_{j=1}^{i-1} w_{ij} (x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta |e_j| + Lh w_{ii} |e_i| + |E_{i,t}(x_i, t, u(t))|, \\ &1 \leq i \leq N. \end{aligned} \quad (16)$$

Let h be so small, that $Lh w_{ii} \leq \frac{1}{2}$, then we easily derive that

$$\begin{cases} e_0 = 0, \\ |e_i| \leq 2Lh \sum_{j=1}^{i-1} (x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta |e_j| + 2|E_{i,t}(x_i, t, u(t))|, \\ 1 \leq i \leq N. \end{cases} \quad (17)$$

Let

$$A = \max_{1 \leq i \leq N} \max_{a \leq t \leq b} |2E_{i,t}(x_i, t, u(t))|,$$

$$B_{ij} = 2Lh(x_i - x_j)^\alpha (\ln |x_i - x_j|)^\beta, \quad \text{for } i > j, \quad -1 < \alpha < 0, \quad \beta = 0 \text{ or } 1, \quad (18)$$

then (17) can be simplified as follows

$$\begin{cases} |e_0| = 0, \\ |e_i| \leq A + \sum_{j=1}^{i-1} B_{ij}|e_j|, \quad 1 \leq i \leq N. \end{cases} \quad (19)$$

Thus the convergence and error estimate of the approximation equation (11) are reduced to estimate $\{|e_i|\}$ satisfying (19). Unfortunately if A and B_{ij} in (19) are defined by (18), then the discrete Gronwall inequality (2) does not hold. In this paper, replacing (2) we shall prove a new generalization of discrete Gronwall inequality.

2. Theorem 1 and its proof

Theorem 1. *If A and B_{ij} are defined by (18) and $\{e_i\}$ satisfies the inequality (19), then there is a positive constant c independent of h , such that*

$$\max_{0 \leq i \leq N} |e_i| \leq ch^{2+\alpha} (\ln h)^\beta. \quad (20)$$

Proof. Since $e_0 = 0$, successively substituting the right side of (19) when $i = n-1, \dots, 1$ into

$$|e_n| \leq A + \sum_{j=1}^{n-1} B_{nj}|e_j|,$$

we derive

$$\begin{aligned} |e_n| &\leq A + A \sum_{j_1=1}^{n-1} B_{nj_1} + A \sum_{j_2=1}^{n-1} B_{nj_2} \sum_{j_1=1}^{j_2-1} B_{j_2j_1} \\ &\quad + \dots + A \sum_{j_{n-1}=1}^{n-1} B_{nj_{n-1}} \sum_{j_{n-2}=1}^{j_{n-1}-1} B_{j_{n-1}j_{n-2}} \dots \sum_{j_1=1}^{j_2-1} B_{j_2j_1}. \end{aligned} \quad (21)$$

In order to estimate $\{|e_n|\}$, we use the following simple inequality: if a nonnegative function $f(x)$ is monotone on $[0, n]$, then

$$\sum_{i=1}^{n-1} f(i) \leq \int_0^n f(x) dx. \quad (22)$$

First we consider the case 1 that $\beta = 0$, i.e., $B_{ij} = 2Lh(x_i - x_j)^\alpha$. Since x^α is monotone and nonnegative, using (22) and setting $x = j_2y$, we get

$$\begin{aligned}
\sum_{j_1=1}^{j_2-1} B_{j_2 j_1} &= 2Lh^{1+\alpha} \sum_{j_1=1}^{j_2-1} (j_2 - j_1)^\alpha \\
&\leq 2Lh^{1+\alpha} \int_0^{j_2} (j_2 - x)^\alpha dx = 2Lh^{1+\alpha} B(1 + \alpha, 1) j_2^{1+\alpha},
\end{aligned} \tag{23}$$

where $B(r, s)$ denote Beta function. Similarly it holds that

$$\begin{aligned}
F_{n,k} &= A \sum_{j_k=1}^{n-1} B_{n j_k} \sum_{j_{k-1}=1}^{j_k-1} B_{j_k j_{k-1}} \cdots \sum_{j_1=1}^{j_2-1} B_{j_2 j_1} \\
&\leq A(2Lh^{1+\alpha})^k n^{k(1+\alpha)} B(1 + \alpha, 1) \cdots B(1 + \alpha, (k-1)(1 + \alpha) + 1).
\end{aligned} \tag{24}$$

However

$$\begin{aligned}
B(1 + \alpha, m(1 + \alpha) + 1) &= \frac{\Gamma(1 + \alpha)\Gamma(m(1 + \alpha) + 1)}{\Gamma((m + 1)(1 + \alpha) + 1)} \\
&= \frac{m}{m + 1} \Gamma(1 + \alpha) \frac{\Gamma(m(1 + \alpha))}{\Gamma((m + 1)(1 + \alpha))}.
\end{aligned} \tag{25}$$

By Stirling formula there exists $\theta_z \in (0, 1)$, such that

$$\Gamma(z) = \sqrt{2\pi} z^{z-\frac{1}{2}} e^{-z} e^{\frac{\theta_z}{12z}}.$$

Letting

$$s = 1 + \alpha,$$

we get that

$$\begin{aligned}
\frac{\Gamma(ms)}{\Gamma((m + 1)s)} &= \left(\frac{m}{m + 1}\right)^{ms-\frac{1}{2}} [(m + 1)s]^{-s} e^s \exp\left(\frac{\theta_m}{12ms} - \frac{\theta_{m+1}}{12(m + 1)s}\right) \\
&\leq e^{\frac{1}{12s}} \left(\frac{e}{s}\right)^s (m + 1)^{-s}.
\end{aligned} \tag{26}$$

Substituting (26) into (25), we obtain

$$B(s, ms + 1) \leq M(m + 1)^{-s}, \tag{27}$$

where $M = \Gamma(s) e^{\frac{1}{12s}} \left(\frac{e}{s}\right)^s$. Applying (27) to (24), we follow that

$$F_{n,k} \leq \frac{A(2L(b-a)^s M)^k}{(k!)^s} \leq A \frac{R^k}{(k!)^s}, \tag{28}$$

where $R = 2L(b-a)sM$. Thus after substituting (28) into (21), we obtain

$$|e_n| \leq A + A \sum_{k=1}^{n-1} \frac{R^k}{(k!)^s} < A \sum_{k=0}^{\infty} \frac{R^k}{(k!)^s} = H A, \tag{29}$$

where

$$H = \sum_{k=0}^{\infty} \frac{R^k}{(k!)^s} < \infty,$$

and H is a positive constant independent of h . However from (9) $A = O(h^{2+\alpha})$, so there is a positive constant c independent of h satisfying

$$\max_{0 \leq i \leq N} |e_i| \leq ch^{2+\alpha}. \quad (30)$$

Secondly consider the case 2 that $\beta = 1$, i.e., $B_{ij} = 2Lh(x_i - x_j)^\alpha (\ln |x_i - x_j|)$. Note that since $\alpha > -1$, we can find such $\varepsilon > 0$, that $\alpha - \varepsilon > -1$. However by the inequality [8]

$$|\ln((i-j)h)| = -\ln((i-j)h) \leq \frac{((i-j)h)^{-\varepsilon}}{\varepsilon e},$$

we derive

$$|B_{ij}| \leq 2Lh[(i-j)h]^\alpha \frac{[(i-j)h]^{-\varepsilon}}{\varepsilon e} = \frac{2L}{\varepsilon e} h[(i-j)h]^{\alpha-\varepsilon}. \quad (31)$$

Setting

$$\alpha_1 = \alpha - \varepsilon, \quad L_1 = \frac{L}{\varepsilon e},$$

and using the results of case 1, we can prove that

$$\max_{0 \leq i \leq N} |e_i| \leq ch^{2+\alpha} (\ln h)^\beta.$$

Thus the proof of Theorem 1 is completed. \square

From Theorem 1 we easy prove the following corollary.

Corollary. If $k(s, t, u)$ satisfies (5), then the solutions $\{u_i\}$ of (11) converges to $\{u(x_i)\}$ as $h \rightarrow 0$, and there exists a constant c independent of h , such that when h is sufficiently small, the error has the estimate

$$\max_{0 \leq i \leq N} |e_i| \leq ch^{2+\alpha} (\ln h)^\beta. \quad (32)$$

3. A numerical example

Considering the second weakly singular Volterra equation

$$x(s) = \frac{11}{18}s^3 - \frac{1}{3}s^3 \ln s + s + \int_0^s \ln |s-t| x^2(t) dt.$$

We know the exact solution is $x(s) = s$, and taking the step width $h = 0.1, 0.05$, the results are listed in the following Table 1.

From the ratio of comparison in Table 1 we can know that the approximation has the accuracy order $O(h^2 |\ln h|)$.

Table 1

t	0.7	0.8	0.9	1.0
$h = 0.1_error$	2.78E-3	2.93E-3	3.29E-3	3.22E-3
$h = 0.05_error$	7.99E-4	8.25E-4	8.45E-4	8.61E-4
ratio of comparison	0.284	0.281	0.257	0.267

Remark. After Gronwall's work [5], many authors generalized Gronwall's inequality and its discrete analogue (see [1,2,8,10,11]). The discrete analogues of the generalizations of Gronwall's inequality are often applied to the numerical treatment of differential equations and integral equations (see [9]), but the numerical treatment of weakly singular integral equations seems to be difficult.

Acknowledgment

The work was supported by the National Natural Science Foundation of China, No. 10171073.

References

- [1] E.F. Beckenbach, D.R. Bellman, *Inequalities*, Springer, Berlin, 1961.
- [2] P.R. Bessack, Comparison theorems and integral inequalities for Volterra integral equations, *Proc. Amer. Math. Soc.* 20 (1969) 61–66.
- [3] L.M. Delves, J. Mohaned, *Computational Methods for Integral Equations*, Cambridge Univ. Press, 1985.
- [4] P.J. Davis, P. Rabinowitz, *Methods of Numeric Integration*, Academic Press, New York, 1984.
- [5] T.H. Gronwall, Note on the derivatives with respect to a parameter of the solutions of a system of differential equations, *Ann. of Math.* 20 (1918) 292–296.
- [6] J.N. Lyness, B.W. Ninham, Numerical quadrature and asymptotic expansions, *Math. Comp.* 21 (1967) 162–178.
- [7] J. Navot, A further extension of Euler–Maclaurin summation formula, *J. Math. Phys.* 41 (1962) 155–184.
- [8] D.S. Mitrinovic, P.M. Vasic, *Analysis Inequalities*, Springer, Berlin, 1970.
- [9] G.I. Marchuk, V.V. Shaidurov, *Difference Methods and Their Extrapolations*, Springer, New York, 1983.
- [10] B.G. Pachpatte, On a certain inequality in the theory of differential equations, *J. Math. Anal. Appl.* 182 (1994) 143–157.
- [11] D. Willett, J.S.W. Wong, On discrete analogues of some generalizations of Gronwall's inequality, *Monatsh. Math.* 69 (1965) 362–367.